

Hallucinogens

What are hallucinogens?

Hallucinogens (also known as 'psychedelics') can make a person see, hear, smell, feel or taste things that aren't really there or are different from how they are in reality.

Some plants such as magic mushrooms can cause hallucinations. Hallucinogens such as LSD can also be made in a lab.

Types of hallucinogens

LSD (Lysergic acid diethylamide)

Also known as acid, trips, tabs, microdots, dots.

In its pure state, LSD is a white odourless powder. However, it usually comes in squares of gelatine or blotting paper that have been dipped or soaked in LSD. LSD is also sometimes sold as a liquid, in a tablet or in capsules.

LSD is usually swallowed, but it can also be sniffed, injected or smoked.

Magic mushrooms

Also known as shrooms, mushies, blue meanies, golden tops, liberty caps.

There are many different types of magic mushrooms. The most common ones in Australia are called golden tops, blue meanies and liberty caps. Magic mushrooms look similar to poisonous mushrooms that can cause a person to become very sick and can result in death.

Magic mushrooms are usually sold as dried mushrooms, a powder or as capsules.

Mushrooms are often eaten fresh, cooked or brewed into a tea. They are sometimes mixed with tobacco or cannabis, and smoked.

Mescaline (peyote cactus)

Also known as cactus, cactus buttons, cactus joint, mesc, mescal.

Mescaline is the active ingredient of the peyote cactus plant. It is also known to be made synthetically in a lab.

In its pure form, mescaline sulphate is a white crystal-like powder. Synthetic mescaline can come in different colours. The peyote cactus contains 'buttons' that can be cut from the root of the plant, and then dried before eating or smoking them.

Effects of hallucinogens

There is no safe level of drug use. Use of any drug always carries some risk. It's important to be careful when taking any type of drug.

Magic mushrooms look similar to poisonous mushrooms that can cause a person to become very sick and can result in death.

Hallucinogens affect everyone differently, based on:

- Size, weight and health
- Whether the person is used to taking it
- Whether other drugs are taken around the same time
- The amount taken
- The strength of the drug (varies from batch to batch)

The effects of hallucinogens can last for 4 to 12 hours and can be different depending on which type of hallucinogen is used. The following may be experienced during this time:

- Feeling happy and relaxed
- Seeing and hearing things that aren't there
- Confusion and trouble concentrating
- Dizziness
- Blurred vision
- Clumsiness
- Fast or irregular heart beat
- Breathing quickly
- Vomiting
- Sweating and chills
- Numbness

Bad trips

Sometimes a 'bad trip' may be experienced, involving a disturbing hallucination. This can lead to panic and risky behaviour, like running across a road or attempting suicide.

If a large amount or a strong batch is taken the negative effects of hallucinogens are more likely.

Coming down

In the following days after using hallucinogens, the following may be experienced:

- Anxiety
- Panic attacks
- Depression

Long term effects

People who regularly use hallucinogens may eventually experience flashbacks. Flashbacks are hallucinations that occur weeks, months or even years after the drug was last taken. This can be disturbing, especially when the hallucination is frightening.

Flashbacks can be brought on by using other drugs, stress, tiredness or exercise and usually last for a minute or two.

In addition to flashbacks, regular use of hallucinogens may eventually cause:

- Psychological dependence on hallucinogens
- Financial, work and social problems

Sometimes a 'bad trip' may be experienced, involving a disturbing hallucination.

1 g]b[\U`i W]bc[Ybg`k]h`ch Yf`Xfi [g`

The effects of taking hallucinogens with other drugs – including over-the-counter or prescribed medications – can be unpredictable and dangerous, and could cause:

<U`i W]bc[Ybg`Z`]WZ`gdYYX`cf`YVg]Ugm enormous strain on the heart and body, which can lead to stroke.

<U`i W]bc[Ybg`Z` U`Vt`c`Z`W]bbUV]g`cf`VYbncX]UnYd]bYg. increased clumsiness and chance of vomiting.

K]h`XfUk U`

Psychological withdrawal symptoms are more common than physical symptoms, but as hallucinogens are a range of different drugs, it's not possible to be specific about withdrawal symptoms. People withdrawing from hallucinogens may experience:

- Cravings
- Fatigue
- Irritability
- Reduced ability to experience pleasure

: i fh`Yf`=bZcfa Uh]cb`

The Druginfo website has information on:

- Help & support services
- Treatment
- Withdrawal
- References for this fact sheet

; c`hc`Xfi []bZc`UXZ`cf["Ui`

These fact sheets
are funded by



Factsheet

GhUm]bZcfa YX`

Hk]h`Yf`Vt`a #5i gh8fi [`]b`YX]b`Vt`a #Vt`a dUbm#Ui ghfU]Ub!Xfi [!Zci bXUh]cb` ZUWVcc_`Vt`a #5i ghfU]Ub8fi [: ci bXUh]cb` [fc[k UHW`UXZ`cf["Ui` UXZ`cf["Ui` #gi VgW]VY`

Level 12, 607 Bourke Street, Melbourne
PO Box 818 North Melbourne Victoria Australia 3051

Email [Xfi \[\]bZc`4UXZ`cf\["Ui`](mailto:Xfi []bZc`4UXZ`cf[) Telephone % \$\$` ,) ,) ; (`

© The Australian Drug Foundation, 2014. ABN 66 057 731 192. This fact sheet offers general information only. The Australian Drug Foundation recommends that you obtain professional advice based on your individual circumstances before acting on information in this fact sheet.

<U`i W]bc[Ybg`

'DU[Y` #`

'A Um&\$%`